

Regimen Reference Order – GAST – mod FOLFOX-6

ARIA: GAST – [mod FOLFOX-6 (ADJ)]

Planned Course: Every 14 days for 8 to 12 cycles

Indication for Use: Colorectal Cancer Adjuvant

CVAD: Required (Ambulatory Pump)

Proceed with treatment if:

ANC equal to or greater than $1.5 \times 10^9/L$ AND Platelets equal to or greater than $100 \times 10^9/L$

❖ Contact Physician if parameters not met

SEQUENCE OF MEDICATION ADMINISTRATION

Pre-treatment Requirements

Drug	Dose	CCMB Administration Guideline
Not Applicable		

Treatment Regimen – GAST – mod FOLFOX-6

Establish primary solution 500 mL of: D5W

Drug	Dose	CCMB Administration Guideline
ondansetron	16 mg	Orally 30 minutes pre-chemotherapy
dexamethasone	12 mg	Orally 30 minutes pre-chemotherapy
oxaliplatin	85 mg/m^2	IV in 500 mL D5W over 2 hours oxaliplatin can be infused at the same time as leucovorin through a Y site
leucovorin	400 mg/m^2	IV in 500 mL D5W over 2 hours
fluorouracil	400 mg/m^2	IV Push over 5 minutes
fluorouracil	2400 mg/m^2	IV in D5W continuously over 46 hours by ambulatory infusion device

All doses will be automatically rounded that fall within the DSG Approved Dose Bands. See GAST DSG – Dose Banding document for more information

In the event of an infusion-related hypersensitivity reaction, refer to the ‘Hypersensitivity Reaction Standing Order’

REQUIRED MONITORING

All Cycles

- CBC and biochemistry as per Physician Orders

Recommended Support Medications

Drug	Dose	CCMB Administration Guideline
dexamethasone	8 mg	Orally once daily on Days 2 and 3
prochlorperazine	10 mg	Orally every 6 hours as needed for nausea and vomiting

DISCHARGE INSTRUCTIONS

- Instruct patient to continue taking anti-emetic(s) at home
- Ensure patient has received a home chemotherapy spill kit and instructions for use
- Reinforce applicable safe handling precautions of medications, blood and body fluids for 48 hours after completion of chemotherapy

ADDITIONAL INFORMATION

- oxaliplatin causes cold intolerance and laryngopharyngeal dysesthesia
 - no ice chips or cold drinks
- oxaliplatin may cause progressive, irreversible neuropathy
 - dose modification may be required