

## Regimen Reference Order

### GENU – pembrolizumab (every 21 days) + aXitinib

ARIA: GENU - [pembro q 21 days + aXitinib]

**Planned Course:** pembrolizumab every 21 days up to a maximum of 2 years (35 cycles) AND aXitinib twice daily until disease progression or unacceptable toxicity

**Indication for Use:** Advanced Renal Cell Carcinoma

**Drug Alert:** Immune Checkpoint Inhibitor (pembrolizumab)

**CVAD:** At Provider's Discretion

**Proceed with treatment if:**

***pembrolizumab + aXitinib***

- *ANC equal to or greater than  $1.5 \times 10^9/L$  AND Platelets equal to or greater than  $50 \times 10^9/L$*
- *AST/ALT equal to or less than 3 times the upper limit of normal*
- *Total bilirubin equal to or less than 1.5 times the upper limit of normal*
- *Creatinine clearance equal to or greater than 30 mL/minute*

***aXitinib Maintenance***

- *ANC equal to or greater than  $0.5 \times 10^9/L$  AND Platelets equal to or greater than  $25 \times 10^9/L$*
- *AST/ALT equal to or less than 3 times the upper limit of normal*
- *Total bilirubin equal to or less than 1.5 times the upper limit of normal*
- *Creatinine clearance equal to or greater than 30 mL/minute*

❖ **Contact Physician if parameters not met**

### SEQUENCE OF MEDICATION ADMINISTRATION

#### Pre-treatment Requirements

Drug	Dose	CCMB Administration Guideline
Not Applicable		

#### Treatment Regimen – GENU – pembrolizumab (every 21 days) + aXitinib

Establish primary solution 500 mL of: normal saline

Drug	Dose	CCMB Administration Guideline
<b>pembrolizumab + aXitinib (Cycles 1 to 35)</b>		
aXitinib	5 mg*	Orally twice daily on <b>Days 1 to 21</b> Take with or without food. Swallow whole <b>(Self-administered at home)</b>
<b>Day 1</b>		
pembrolizumab	2 mg/kg	IV in normal saline 50 mL over 30 minutes <i>Use 0.2 or 0.22 micron filter</i>

aXitinib maintenance (Cycle 36 and Onwards)		
aXitinib	5 mg*	Orally twice daily on <b>Days 1 to 21</b> Take with or without food. Swallow whole <b>(Self-administered at home)</b>
<b>Maximum pembrolizumab dose is 200 mg</b> All doses will be automatically rounded that fall within CCMB Approved Dose Bands. See Dose Banding document for more information		
* At the discretion of the Medical Oncologist, aXitinib dose may be increased to 7 mg twice daily and subsequently to 10 mg twice daily if the patient is tolerating therapy OR dose may be reduced to 3 mg twice daily and subsequently to 2 mg twice daily if the patient is experiencing adverse drug reactions		
<b>aXitinib (INLYTA®) available dosage strengths: 1 mg and 5 mg tablets</b> <b>Classification: Cytotoxic, Hazardous</b>		

In the event of an infusion-related hypersensitivity reaction, refer to the ‘Hypersensitivity Reaction Standing Order’

## REQUIRED MONITORING

### Throughout treatment (aXitinib)

- Blood pressure
  - At baseline
  - After 1 week
  - Frequently thereafter (at least monthly)
- Urine protein
  - Urinalysis for protein: Where urinalysis is not possible, use dipstick. If lab urinalysis for protein is greater than or equal to 1 g/L or dipstick proteinuria shows 2+ or 3+, notify prescriber
  - At baseline
  - Every 3 to 4 months or as clinically indicated

### All Cycles

- CBC, serum creatinine, urea, electrolytes, liver enzymes, total bilirubin, albumin, glucose and TSH as per Physician Orders

### Cycles 1 to 35 (pembrolizumab)

- Medical oncologist or designate (i.e. family practitioner in oncology) must assess patient for immune-mediated adverse reactions prior to each cycle
- Full vital signs (temperature, heart rate, respiratory rate, blood pressure and O<sub>2</sub> saturation) at baseline and as clinically indicated
- No observation period is required after pembrolizumab administration. Patient can be discharged from treatment room if stable whether they had a reaction or not

## Recommended Support Medications

Drug	Dose	CCMB Administration Guideline
None required		

---

## DISCHARGE INSTRUCTIONS

---

### All Cycles

- Patient should monitor blood pressure at home and record measurements on blood pressure log. This should be done daily for at least the first 2 cycles
- Contact your cancer care team if systolic blood pressure is greater than or equal to 170 mmHg or diastolic blood pressure is greater than 95 mmHg on two consecutive readings
- aXitinib has potential for significant drug-drug interactions. Patients should notify clinic prior to starting any new medication
- Avoid grapefruit and grapefruit juice, Seville oranges (i.e. orange marmalade), and starfruit
- Reinforce applicable safe handling precautions of medications, blood and body fluids while on aXitinib

### Cycles 1 to 35 (pembrolizumab)

- Patient should be instructed to contact their cancer team immediately if symptoms of hypersensitivity reactions occur after discharge
- Confirm that patient has received the CCMB Immune Checkpoint Inhibitor Medical Alert wallet card
- Reinforce to patient the immune-mediated adverse reactions and importance of reporting immediately
  - For severe symptoms, the patient should be instructed to go to the nearest emergency room. Oncologist on call should be contacted

---

## ADDITIONAL INFORMATION

---

- pembrolizumab is an Immune Checkpoint Inhibitor. Consult with oncologist for immune-mediated adverse reactions; corticosteroids are often indicated
- aXitinib can increase risk of hypertension, hemorrhage, wound healing complications and thromboembolic events
- aXitinib can cause rare but serious side effects such as GI perforation and fistulas, reversible posterior leukoencephalopathy syndrome (RPLS) and cardiac failure