

Regimen Reference Order – H&N – nivolumab

ARIA: H&N – [nivolumab q 14 days]

H&N – [nivolumab q 28 days]

Planned Course: Every 14 days until disease progression or unacceptable toxicity
OR
Every 28 days until disease progression or unacceptable toxicity

Indication for Use: Squamous Cell Cancer Head and Neck Metastatic

Drug Alert: Immune Checkpoint Inhibitor

CVAD: At Provider's Discretion

Proceed with treatment if:

- **ANC equal to or greater than $1.5 \times 10^9/L$ AND Platelets equal to or greater than $50 \times 10^9/L$**
- **AST/ALT less than 3 times upper limit of normal**
- **Total bilirubin less than 1.5 times upper limit normal**
- **Creatinine clearance greater than 30 mL/min**
- ❖ **Contact Physician if parameters not met**

SEQUENCE OF MEDICATION ADMINISTRATION

Pre-treatment Requirements

Drug	Dose	CCMB Administration Guideline
Not Applicable		

Treatment Regimen – H&N - nivolumab

Establish primary solution 500 mL of: normal saline

Drug	Dose	CCMB Administration Guideline
nivolumab	3 mg/kg (every 14 days) OR	IV in normal saline 100 mL over 30 minutes Use 0.2 or 0.22 micron filter <i>Concentration dependent drug: Pharmacy will adjust diluent volume to ensure drug stability</i>
	6 mg/kg (every 28 days)	IV in normal saline 100 mL over 30 minutes Use 0.2 or 0.22 micron filter <i>Concentration dependent drug: Pharmacy will adjust diluent volume to ensure drug stability</i>

Maximum nivolumab dose is 240 mg (every 14 days) OR 480 mg (every 28 days)

All doses will be automatically rounded that fall within the DSG Approved Dose Bands. See H&N DSG – Dose Banding document for more information

Flush after each medication:

- 50 mL over 6 minutes (500 mL/hr)

In the event of an infusion-related hypersensitivity reaction, refer to the 'Hypersensitivity Reaction Standing Order'

REQUIRED MONITORING

All Cycles

- CBC, serum creatinine, urea, electrolytes, AST, ALT, total and direct bilirubin, TSH and glucose as per Physician Orders
- Medical oncologist or designate (i.e. family practitioner in oncology) must assess patient for immune-mediated adverse reactions prior to each cycle
- Full vital signs (temperature, heart rate, respiratory rate, blood pressure and O₂ saturation) at baseline and as clinically indicated
- No observation period is required
- Patient can be discharged from treatment room if stable whether they had a reaction or not

Recommended Support Medications

Drug	Dose	CCMB Administration Guideline
Not Applicable		

DISCHARGE INSTRUCTIONS

- Confirm that patient has received the CCMB Immune Checkpoint Inhibitor Medical Alert wallet card
- Reinforce to patient the immune-mediated adverse reactions and importance of reporting immediately
 - For severe symptoms, the patient should be instructed to go to the nearest emergency room. Oncologist on call should be contacted

ADDITIONAL INFORMATION

- nivolumab is an Immune Checkpoint Inhibitor. Consult with oncologist for immune-mediated adverse reactions; corticosteroids are often indicated