

## Regimen Reference Order – CLL – R-CD

ARIA: CLL – [R-CD]

Planned Course: Every 28 days for 6 cycles

Indication for Use: Autoimmune hemolytic anemia

CVAD: At Provider's Discretion

### Proceed with treatment if:

#### Cycle 1

- Proceed with treatment regardless of CBC

#### Cycles 2 to 6

- ANC and platelets are the same or greater than pre-treatment value (*prior to Cycle 1, Day 1*)
  - ❖ Contact Hematologist if parameters not met

**Note:** Hepatitis B serology results must be reviewed in accordance with CCMB Policy Hepatitis B Monitoring for Oncology and Hematology Patients

## SEQUENCE OF MEDICATION ADMINISTRATION

### Pre-treatment Requirements

Drug	Dose	CCMB Administration Guideline
Not Applicable		

### Treatment Regimen – CLL – R-CD

Establish primary solution 500 mL of: normal saline

Drug	Dose	CCMB Administration Guideline
<b>Cycle 1</b>		
<b>Day 1</b>		
cetirizine	10 mg	Orally 30 minutes prior to riTUXimab
acetaminophen	650 mg	Orally 30 minutes prior to riTUXimab
dexamethasone	40 mg	IV in normal saline 50 mL over 15 minutes
<b>Wait 30 minutes after completion of IV pre-medications before starting riTUXimab</b>		
riTUXimab (IV brand name specific)	375 mg/m <sup>2</sup>	<p><b>Slow infusion</b> (if greater than 6 months since last riTUXimab dose or no previous riTUXimab): IV made up to a final concentration of 1 mg/mL in normal saline. Start at 50 mg/hr for 60 minutes and escalate infusion rate in 50 mg/hr increments every 30 minutes to maximum of 400 mg/hr</p> <p><i>*Nursing Alert: IV tubing is primed with riTUXimab</i></p> <p><i>*Alert: Ensure brand name on prescription label (indicated in brackets on prescription label) matches prescribed order</i></p> <p><i>*Alert: Pharmacy to ensure final volume on label</i></p> <p><b>OR</b></p>

		<p><b>Slow infusion (if equal to or less than 6 months since last riTUXimab dose):</b> IV made up to a final concentration of 1 mg/mL in normal saline. Start at 100 mg/hr for 30 minutes and escalate infusion rate in 100 mg/hr increments every 30 minutes to maximum of 400 mg/hr</p> <p><i>*Nursing Alert: IV tubing is primed with riTUXimab</i></p> <p><i>*Alert: Ensure brand name on prescription label (indicated in brackets on prescription label) matches prescribed order</i></p> <p><i>*Alert: Pharmacy to ensure final volume on label</i></p>
ondansetron	16 mg	Orally 30 minutes pre-chemotherapy
cyclophosphamide	750 mg/m <sup>2</sup>	IV in normal saline 250 mL over 1 hour
<b>Days 2 to 7</b>		
dexamethasone	12 mg	Orally once daily in the morning with food (Self-administered at home)
<b>Cycle 2 onwards</b>		
<b>Day 1</b>		
cetirizine	10 mg	Orally 30 minutes prior to riTUXimab
acetaminophen	650 mg	Orally 30 minutes prior to riTUXimab
dexamethasone	20 mg	IV in normal saline 50 mL over 15 minutes
<b>Wait 30 minutes after completion of IV pre-medications before starting riTUXimab</b>		
riTUXimab (Subcutaneous)	1400 mg (1400 mg = 11.7 mL)	<p><b>Subcutaneous:</b> Administer over 5 minutes into abdomen Syringe should be held in hand for 5 minutes to warm up and decrease viscosity Use 25G needle</p> <p><i>*Nursing Alert: Ensure subcutaneous riTUXimab formulation is used (riTUXimab-hyaluronidase, human)</i></p>
<b>OR</b>		
riTUXimab (IV brand name specific)	375 mg/m <sup>2</sup>	<p><b>Slow infusion:</b> IV made up to a final concentration of 1 mg/mL in normal saline. Start at 100 mg/hr for 30 minutes and escalate infusion rate in 100 mg/hr increments every 30 minutes to maximum of 400 mg/hr</p> <p><i>*Nursing Alert: IV tubing is primed with riTUXimab</i></p> <p><i>*Alert: Ensure brand name on prescription label (indicated in brackets on prescription label) matches prescribed order</i></p> <p><i>*Alert: Pharmacy to ensure final volume on label</i></p> <p><b>OR</b></p> <p><b>Rapid infusion:</b> IV in normal saline over 90 minutes: Infuse 50 mL of a 250 mL bag (or 100 mL of a 500 mL bag) over 30 minutes, then infuse the remaining 200 mL (or 400 mL of a 500 mL bag) over 60 minutes</p> <p><i>*Alert: Ensure brand name on prescription label (indicated in brackets on prescription label) matches prescribed order</i></p> <p><i>*Alert: Pharmacy to ensure final volume on label</i></p>

ondansetron	16 mg	Orally 30 minutes pre-chemotherapy
cyclophosphamide	750 mg/m <sup>2</sup>	IV in normal saline 250 mL over 1 hour
<b>Days 2 to 7</b>		
dexamethasone	12 mg	Orally once daily in the morning with food <b>(Self-administered at home)</b>
All doses will be automatically rounded that fall within CCMB Approved Dose Bands. See Dose Banding document for more information		

**In the event of an infusion-related hypersensitivity reaction, refer to the 'Hypersensitivity Reaction Standing Order'**

## REQUIRED MONITORING

### Hepatitis B serology

- Hepatitis B surface antigen and Hepatitis B core antibody (drawn within preceding 5 years)

### All Cycles

#### Day 1

- CBC, serum creatinine, urea, liver enzymes, electrolytes, LDH, total bilirubin, uric acid and albumin as per Physician Orders

### INTRAVENOUS ritUXimab

- Full vital signs (temperature, heart rate, respiratory rate, blood pressure and O<sub>2</sub> saturation) at baseline and as clinically indicated
- No observation period is required after ritUXimab administration. Patient can be discharged from treatment room if stable whether they had a reaction or not

### SUBCUTANEOUS ritUXimab

- Full vital signs (temperature, heart rate, respiratory rate, blood pressure and O<sub>2</sub> saturation) at baseline, at discharge, and as clinically indicated
- 15-minute observation period required after each dose**

## Recommended Support Medications

Drug	Dose	CCMB Administration Guideline
sulfamethoxazole-trimethoprim	800/160mg	Orally twice daily on Saturdays and Sundays only
metoclopramide	10 – 20 mg	Orally every 4 hours as needed for nausea and vomiting

---

## DISCHARGE INSTRUCTIONS

---

- Patients should be instructed to contact their cancer team immediately if symptoms of hypersensitivity reactions occur after discharge
  - Instruct patient to:
    - Continue taking anti-emetic(s) at home
    - Maintain oral intake of 2000 mL (8 glasses) of fluid daily at home
    - Empty bladder every 2 hours while awake and at bedtime for 24 hours
    - Obtain immediate assistance as per your clinic's contact instructions if:
      - Symptoms of hemorrhagic cystitis (e.g. dysuria, hematuria)
      - Unable to drink recommended amount of fluid
  - sulfamethoxazole-trimethoprim is prescribed for *Pneumocystis jirovecii* pneumonia prophylaxis. Remind patient to take sulfamethoxazole-trimethoprim at home
  - dexamethasone is a cancer therapy in this treatment regimen. Remind patient to take dexamethasone at home
  - Reinforce applicable safe handling precautions of medications, blood and body fluids for 48 hours after completion of chemotherapy
- 

## ADDITIONAL INFORMATION

---

- Due to the risk of reactivation of Hepatitis B virus (HBV) while on this treatment regimen, prescriber must adhere to CCMB Policy ***Hepatitis B Monitoring for Oncology and Hematology Patients*** for ordering and interpreting HBV serology and prescribing antiviral prophylaxis
- Administering nurse must document any infusion-related reactions with any dose of riTUXimab
- Ensure there were **no Grade 3 or 4** infusion-related reactions with any previous dose prior to administering riTUXimab via Rapid Infusion or Subcutaneous injection
- Intravenous riTUXimab formulation is available from more than one manufacturer and uses several different brand names. Brand name will be indicated in brackets after riTUXimab. **Ensure prescription label matches the brand name on prescribed order for intravenous riTUXimab**