

Regimen Reference Order – THOR – CAV

ARIA: LUNG – [CAV]

Planned Course: Every 21 days for 4 to 6 cycles

Indication for Use: Small Cell Lung Cancer

CVAD: Preferred (VESICANT INVOLVED)

Proceed with treatment if:

ANC equal to or greater than $1.5 \times 10^9/L$ AND Platelets equal to or greater than $100 \times 10^9/L$

❖ Contact Physician if parameters not met

Note: Hepatitis B serology results must be reviewed in accordance with CCMB Policy Hepatitis B Monitoring for Oncology and Hematology Patients

SEQUENCE OF MEDICATION ADMINISTRATION

Pre-treatment Requirements

Drug	Dose	CCMB Administration Guideline
Not Applicable		

Treatment Regimen – THOR – CAV

Establish primary solution 500 mL of: normal saline

Drug	Dose	CCMB Administration Guideline
Day 1		
ondansetron	16 mg	Orally 30 minutes pre-chemotherapy
dexamethasone	12 mg	Orally 30 minutes pre-chemotherapy
DOXOrubicin	50 mg/m^2	IV push over 10 to 15 minutes
vinCRISTine	1.2 mg/m^2 ; maximum dose 2 mg	IV in normal saline 25 mL over 2 to 3 minutes by gravity infusion
cyclophosphamide	1000 mg/m^2	IV in normal saline 250 mL NS over 1 hour

All doses will be automatically rounded that fall within CCMB Approved Dose Bands. See Dose Banding document for more information

In the event of an infusion-related hypersensitivity reaction, refer to the 'Hypersensitivity Reaction Standing Order'

REQUIRED MONITORING

Hepatitis B serology

- Hepatitis B surface antigen and Hepatitis B core antibody (drawn within preceding 5 years)

Cardiac monitoring

- Left Ventricular Ejection Fraction (LVEF) monitoring is recommended at baseline and as clinically indicated

All Cycles

- CBC, serum creatinine, urea, liver enzymes and electrolytes as per Physician Orders

Recommended Support Medications

Drug	Dose	CCMB Administration Guideline
dexamethasone	8 mg	Orally once daily on Days 2 and 3
metoclopramide	10 – 20 mg	Orally every 4 hours as needed for nausea and vomiting

DISCHARGE INSTRUCTIONS

- Instruct patient to:
 - Continue taking anti-emetic(s) at home
 - Maintain oral intake of 2000 mL (8 glasses) of fluid daily at home
 - Empty bladder every 2 hours while awake and at bedtime for 24 hours
 - Obtain immediate assistance as per your clinic's contact instructions if:
 - Symptoms of hemorrhagic cystitis (e.g. dysuria, hematuria)
 - Unable to drink recommended amount of fluid
- Reinforce applicable safe handling precautions of medications, blood and body fluids for 48 hours after completion of chemotherapy

ADDITIONAL INFORMATION

- Due to the risk of reactivation of Hepatitis B virus (HBV) while on this treatment regimen, prescriber must adhere to CCMB Policy **Hepatitis B Monitoring for Oncology and Hematology Patients** for ordering and interpreting HBV serology and prescribing antiviral prophylaxis