

Regimen Reference Order

THOR – gemcitabine + CISplatin (Adjuvant)

ARIA: LUNG – [gemcitabine + CIS (ADJ)]

Planned Course: Every 21 days for 4 cycles
Indication for Use: Lung Cancer Non-Small Cell Adjuvant

CVAD: At Provider's Discretion

Proceed with treatment if:

Day 1

- ANC equal to or greater than $1.5 \times 10^9/L$ AND Platelets equal to or greater than $100 \times 10^9/L$
- Creatinine clearance greater than 45 mL/minute

Day 8

- ANC equal to or greater than $1.5 \times 10^9/L$ AND Platelets equal to or greater than $100 \times 10^9/L$
- ❖ Contact Physician if parameters not met

SEQUENCE OF MEDICATION ADMINISTRATION

Pre-treatment Requirements

Drug	Dose	CCMB Administration Guideline
Not Applicable		

Treatment Regimen – THOR – gemcitabine + CISplatin (Adjuvant)

Establish primary solution 500 mL of: normal saline

Drug	Dose	CCMB Administration Guideline
Day 1		
magnesium sulfate	2 g	IV in normal saline 1000 mL over 2 hours (Pre hydration)
aprepitant	125 mg	Orally 1 hour pre-chemotherapy
ondansetron	16 mg	Orally 30 minutes pre-chemotherapy
dexamethasone	12 mg	Orally 30 minutes pre-chemotherapy
OLANzapine	2.5 mg	Orally 30 minutes pre-chemotherapy
gemcitabine	1000 mg/m^2	IV in normal saline 250 mL over 30 minutes
CISplatin	75 mg/m^2	IV in normal saline 500 mL over 1 hour <i>*Alert: CISplatin infusion must be complete prior to mannitol administration</i>
mannitol	12.5 g	IV in normal saline 1000 mL over 2 hours (Post hydration)

Day 8		
dexamethasone	8 mg	Orally 30 minutes pre-chemotherapy
gemcitabine	1000 mg/m ²	IV in normal saline 250 mL over 30 minutes

In the event of an infusion-related hypersensitivity reaction, refer to the 'Hypersensitivity Reaction Standing Order'

REQUIRED MONITORING

All Cycles

Day 1

- CBC, serum creatinine, urea, electrolytes, liver enzymes and total bilirubin as per Physician Orders
- Baseline blood pressure prior to magnesium infusion and repeat 15 minutes after start of magnesium infusion

Day 8

- CBC

Recommended Support Medications

Drug	Dose	CCMB Administration Guideline
aprepitant	80 mg	Orally once daily on Days 2 and 3
dexamethasone	8 mg	Orally once daily on Days 2 to 4
OLANzapine	2.5 mg	Orally the evening of Day 1 then twice daily on Days 2, 3 and 4. Also use OLANzapine 2.5 to 5 mg AS NEEDED for breakthrough nausea and vomiting (including Days 1 to 4) up to a maximum of 10 mg per day. Contact clinic if nausea/vomiting is not adequately controlled

DISCHARGE INSTRUCTIONS

- Instruct patient to continue taking anti-emetic(s) at home
- Reinforce applicable safe handling precautions of medications, blood and body fluids for 48 hours after completion of chemotherapy

ADDITIONAL INFORMATION

- CISplatin is ototoxic and nephrotoxic
- CISplatin can cause hypomagnesemia