

## COLONCHECK RESOURCE REFERENCE LISTING

---

- Hewitson P, Glasziou P, Irwig L & Watson E (2007). Screening for colorectal cancer using the fecal occult blood test, Hemoccult. *Cochrane Database of Systemic Review*, 1.
- Leddin, D, Lieberman, D. et al. Clinical practice guideline on screening for colorectal cancer in individuals with a family history of nonhereditary colorectal cancer or adenoma: The Canadian Association of Gastroenterology Banff consensus. *Can J Gastroenterology* 2018;155:1325-1347.
- Leddin D, Enns R et al. Canadian Association of Gastroenterology position statement on screening individuals at average risk for developing colorectal cancer,; *Can J Gastroenterology* 2010;24(12):705-14.
- Leddin D, Enns R, Hilsden R, Fallone CA, Rabeneck L, Sadowski DC, Singh H. Colorectal cancer surveillance after index colonoscopy: Guidance from the Canadian Association of Gastroenterology. *Can J Gastroenterology* 2013;27(4):224-228.
- Lieberman DA, Rex DK, Winawer SJ, Giardiello FM, Johnson DA, Levin TR. Guidelines for colonoscopy surveillance after screening and polypectomy: A consensus update by the US Multi-Society Task Force on Colorectal Cancer. *Gastroenterology*. 2012;143:844–57.
- Recommendations on screening for colorectal cancer in primary care; Canadian Task Force on Preventative Health Care. *CMAJ* 2016. doi:10.1503/cmaj.151115
- Singh H, Turner D, Xue L, Targownik LE, Bernstein CN. Risk of developing colorectal cancer following a negative colonoscopy examination: Evidence for a 10-year interval between colonoscopies. *JAMA*. 2006;295(20):2366–2373. doi:10.1001/jama.295.20.2366
- Tinmouth J. et al. Colorectal cancer screening in average risk populations: Evidence summary. *Canadian Journal of Gastroenterology & Hepatology*. Volume 2016, Article ID 2878149, 18 pages. doi.org/10.1155/2016/2878149